

MENINGITIS WAIVER

New York State Department of Health requires each student to indicate meningitis compliance by providing a waiver of the vaccine OR providing medical documentation of date of vaccine. The Student Health Center does NOT provide this vaccine. You may contact your local county health department or your personal physician to obtain this vaccination.

| Student Last Name | First | College ID# | Date of Birth |
|-------------------|-------|-------------|---------------|
|-------------------|-------|-------------|---------------|

WAIVER: I have reviewed the attached fact sheet regarding meningococcal disease. I am fully aware of the risks associated with this disease and of the availability and effectiveness of the vaccine. I have elected NOT to get the vaccine.

Signature of Student (*Parent/Guardian if student under 18*) _____ Date _____

**NEW YORK STATE DEPARTMENT OF HEALTH
Bureau of Communicable Disease Control
Meningococcal Disease**

What is meningococcal disease? Meningococcal disease is a severe bacterial infection of the bloodstream or meninges (a thin lining covering the brain and spinal cord) caused by the meningococcus germ.

Who gets meningococcal disease? Anyone can get meningococcal disease, but it is more common in infants and children. For some adolescents, such as first year college students living in dormitories, there is an increased risk meningococcal disease. Every year in the United States approximately 2, 500 people are infected and 300 die from the disease.

How is the meningococcus germ spread? The meningococcus germ is spread by direct close contact with nose or throat discharges of an infected person.

What are the symptoms? High fever, headache, vomiting, stiff neck, and a rash are symptoms of meningococcal disease. The symptoms may appear 2 to 10 days after exposure, but usually within 5 days. Among people who develop meningococcal disease, 10-15% dies, in spite of treatment with antibiotics. Of those who live, permanent brain damage, hearing loss, kidney failure, and loss of use of arms or legs, or chronic nervous system problems can occur.

What is the treatment for meningococcal disease? Antibiotics, such as penicillin G or ceftriaxone, can be used to treat people with meningococcal disease.

Should people who have been in contact with a diagnosed case of meningococcal meningitis be treated? Only people who have been in close contact (household members, intimate contacts, health care personnel performing mouth-to-mouth resuscitation, day care center playmates, etc.) need to be considered for preventive treatment. Such people are usually advised to obtain a prescription for a special antibiotic (rifampin, ciprofloxacin, or ceftriaxone) from their physician. Casual contact, as might occur in a regular classroom, office, or factory setting, is not usually significant enough to cause concern.

Is there a vaccine to prevent meningococcal meningitis? In February 2005, the CDC recommended a new vaccine, known as Menactra™, for use to prevent meningococcal disease in people 11-55 years of age. The previously licensed version of this vaccine, Menomune™, is available for children 2-10 years old and adults older than 55 years. Both vaccines are 85% to 100% effective in preventing the 4 kinds of meningococcus germ (types A, C, Y, W-135). These 4 types cause about 70% of the disease in the United States. Because the vaccines do not include B, which accounts for about one-third of cases in adolescents, they do not prevent all cases of meningococcal disease.

Are the vaccines safe? Are there adverse side effects to the vaccine? Both vaccines are currently available and both are safe and effective vaccines. However, both vaccines may cause mild and infrequent side effects, such as redness and pain at the injection site lasting up to two days.

Who should get the meningococcal vaccine? The vaccine is recommended for all adolescents entering middle school (11-12 years old) and high school (15 years old), and all first year college students living in dormitories. However, the vaccine will benefit all teenagers and young adults in the United States. Also at increased risk are people with terminal complement deficiencies or asplenia, some laboratory workers, and travelers to endemic areas of the world.

What is the duration of protection from the vaccine? Menomune™, the older vaccine, requires booster doses every 3 to 5 years. Although research is still pending, the new vaccine, Menactra™ will probably not require booster doses.

How do I get more information about meningococcal disease and vaccination? Contact your family physician or your student health service. Additional information is also available on the websites of the NYS DOH, www.health.state.ny.us; the CDC & Prevention, www.cdc.gov/ncid/dbmd/diseaseinfo; and the American College Health Association, www.acha.org.